

Country review and roadmap for action



Papua New Guinea

Cervical cancer is one of the most preventable cancers. However, in 2018, an estimated 311,000 women died from cervical cancer globally.¹ Most of these deaths occurred in low- and lower-middle income countries (LMICs) due to inadequate access to cervical cancer prevention.

In November 2020, the WHO launched a Global Strategy to accelerate the elimination of cervical cancer as a public health problem. The Strategy proposes an elimination threshold of 4 cases per 100,000 women, achieved by implementing the triple intervention targets by 2030:

- 90% of girls fully vaccinated with the HPV vaccine by age 15.
- 70% of women screened with a high-performance test (such as the HPV test) by 35, and again by 45 years.
- 90% of women identified with cervical precancer or cervical cancer receive adequate treatment and care.²

If this Strategy is adopted, **a total of 74 million cervical cancer cases could be prevented and 62 million women's lives could be saved over the next century.**^{3,4} This Strategy also represents the most cost-effective approach across 95% of countries analyzed, and is expected to offer immense economic and societal benefits, with **an estimated US\$ 3.20 returned to the economy, rising to US\$ 26.00 when societal benefits are incorporated, for every dollar invested through 2050** due to increases in women's participation in the workforce.²

In Papua New Guinea (PNG), cervical cancer is the second most common cancer in women, with 1,024 new cases (29.1 per 100,000 women) and 663 deaths (19.8 per 100,000 women) in 2018.¹ The HIV prevalence among adults (15-49 years) was 0.9% and an estimated 62% of people living with HIV were on treatment in 2018.⁵ It was predicted that without any intervention, **a total of 60,232 women in PNG will die from cervical cancer by 2070 and 175,166 by 2120.**³

Pilot studies for cervical cancer screening have been conducted in the country over the past decade;^{6,7} however, there is no national program for either HPV vaccination or cervical cancer screening in Papua New Guinea. To reach the elimination targets, organized national programmes for both HPV vaccination and cervical cancer screening need to be implemented, and competencies of health workforce will need to be strengthened across all three pillars (HPV vaccination, cervical cancer screening and treatment) for long-term success on the path towards elimination.

Papua New Guinea could eliminate cervical cancer by 2085

The introduction of HPV vaccines into the national immunization programme is a crucial first step to be able to achieve a high coverage of HPV vaccination. Scale-up of HPV vaccination coverage could be facilitated by considering multisectoral

delivery platforms, including a school-based strategy targeting adolescent females aged 9-14 years, or innovative community-based strategies. It is not cost-effective to vaccinate boys as well as girls. It is also important to secure sufficient and affordable HPV vaccines, through a concerted effort to negotiate for reasonable prices and with international partnerships and support, including from Gavi. Improving monitoring systems and implementing vaccination registers are critical for ensuring that high coverage is met. Educating the community on the importance of HPV vaccination to prevent cervical cancer and having strategies to deal with misinformation are also important for maintaining confidence in the programme.

If the three pillars of elimination are established, Papua New Guinea could avert over 149,000 deaths due to cervical cancer over the next century.

Cervical cancer screening programmes represent the second pillar of the three-pronged Strategy. These screening programmes could be integrated with existing services such as primary care services, for instance by offering HPV testing at sexual health clinics, ante-natal care consultations, family planning consultations or well women clinics. Integrating cervical cancer screening with facilities that offer services for HIV control should be considered to ensure women at high risk of cervical cancer have access to cervical cancer screening. This would minimize unnecessary referrals and reduce transport costs and waiting time for women. Understanding social and cultural barriers will help create an environment where women feel comfortable attending screening visits.

Additionally, it is essential to provide information and education for women on the importance

of cervical cancer screening, and potentially offering self-collection for HPV testing, or point-of-care HPV testing in rural areas, so that women who need precancer treatment can be treated in the same visit. Screening scale-up should utilize clinically validated high sensitivity tests, and strengthening of laboratory services and quality assurance programmes will ensure the programme is effective. Implementing surveillance and monitoring systems are also critical for ensuring women are not lost to follow-up and for monitoring programme success nationally.


Lastly, developing and implementing national cervical cancer management guidelines is essential to providing women with high quality cancer treatment and care. Establishing strong referral networks

between all aspects of the cancer care spectrum are needed to ensure the timely management of patients and reduce loss-to-follow-up. Diagnosis, treatment, and monitoring of invasive cervical cancer will require high-quality pathology services and oncology services. Safe, effective, and timely surgery accompanied by supportive services such as anesthetic services and intensive care units are important for the treatment of early stage cancers. Furthermore, expanding access to radiotherapy units and access to chemotherapy services, as well as palliative care services, are crucial.

The Elimination Implementation Checklist on the final page summarizes the steps that countries should take to ensure the three pillars for elimination are successfully implemented and maintained over the longer term.

Figure 1:

Contribution of each elimination pillar to deaths averted in Papua New Guinea.



The contribution of vaccination is calculated by comparing model outputs for vaccination versus status quo. The contribution of screening is calculated by comparing modelled outputs for vaccination and screening versus vaccination. The contribution of cancer treatment is calculated by comparing modelled outputs for vaccination, screening and cancer treatment versus vaccination and screening. Contributions of the pillars may not sum to total due to rounding.

Table 1:

Cervical cancer deaths averted, attributable to each pillar of the elimination strategy.

Year	Vaccination	Screening & precancer treatment	Cancer treatment	Total
2020-2030	4	157	936	1,096
2020-2070	18,986	13,892	8,284	41,163
2020-2120	113,510	23,028	12,998	149,537

Prospects for Cervical Cancer Elimination in Papua New Guinea

Concerted action now across all three elimination pillars is critical. The timing of impacts will vary, but action now will have substantial long-term impacts.

If all three elimination pillars are reached by 2030, a total of 1,096 cervical cancer deaths will be prevented by 2030, 41,163 deaths prevented by 2070 and 149,537 cervical cancer deaths by 2120 (Table 1).

PNG could eliminate cervical cancer by 2085.

The effects of HPV vaccination will be seen after a few decades (Figure 1). HPV vaccination would have limited impact before 2030, but the results will be seen in subsequent decades: HPV vaccination could prevent nearly 18,986 cervical cancer deaths by 2070, and 113,510 cervical cancer deaths by 2120 (Table 1).

The effects of cervical cancer screening and precancer treatment will be seen mainly in the mid-term, after 2030 (Figure 1). Cervical cancer screening would prevent an additional 157 cervical cancer deaths by 2030, 13,892 deaths by 2070 and 23,028 cervical cancer deaths by 2120 (Table 1).

Cervical cancer treatment scale-up will have immediate impacts on deaths averted (Figure 1). Scale-up of cancer treatment would prevent an additional 936 cervical cancer deaths by 2030, 8,284 deaths by 2070 and 12,998 cervical cancer deaths by 2120 (Table 1).

Papua New Guinea is at a crucial stage in its efforts to prevent and treat cervical cancer, and a concerted strategic plan to accelerate the elimination of cervical cancer needs to be developed and implemented. There will be many challenges along the way, including vaccine and screening test supply and delivery challenges, human resource capacity and the infrastructure challenges associated with scale-up of invasive cancer diagnostics, treatment, and supportive and palliative care services. However, the tools and technology are now available; research has shown that elimination strategies are cost-effective and will prevent over 149,000 deaths over the next century. Both HPV vaccination and cervical cancer screening and treatment have been identified by WHO as best buys and are included in the list of national health plan recommendations by WHO. The elimination agenda is a component of the United Nations Global Strategy for Women's, Children's and Adolescent's Health. Achieving cervical cancer elimination in Papua New Guinea will support several sustainable development goals (SDGs) and targets, including SDG 3 (good health and well-being), SDG 5 (gender equity) and SDG 10 (reducing inequalities). The implementation of HPV vaccination, cervical cancer screening and cancer treatment access will be facilitated by the full realization of universal health care in Papua New Guinea, and collaboration with other sectors for sharing knowledge, financial resources and expertise in achieving the SDG goals will be crucial. Cervical cancer elimination is within our reach and we urge all stakeholders to support this effort by investing in the comprehensive approach as a national priority.

References

1. Ferlay J, Ervik M, Lam F, et al. Global Cancer Observatory: Cancer Today [Internet]. Lyon, France: International Agency for Research on Cancer; 2018 [cited 2020 October 10]. Available from: <https://gco.iarc.fr/today>.
2. Global strategy to accelerate the elimination of cervical cancer as a public health problem. Geneva: World Health Organization; 2020. Licence: CC BY-NC-SA 3.0 IGO.
3. Canfell K, Kim J J, Brisson M, et al. Mortality impact of achieving WHO cervical cancer elimination targets: a comparative modelling analysis in 78 low-income and lower-middle-income countries. *Lancet* 2020;395:591-603.
4. Brisson M, Kim J J, Canfell K, et al. Impact of HPV vaccination and cervical screening on cervical cancer elimination: a comparative modelling analysis in 78 low-income and lower-middle-income countries. *Lancet* 2020;395:575-90.
5. UNAIDS. Country factsheets [Internet]. UNAIDS; 2018 [cited 2020 December 3]. Available from: <https://www.unaids.org/en/regionscountries/countries>.
6. Mola G., Posanai E., Augerea L., et al. Final report of the Ministerial Task Force on Prevention and Treatment of Cancer of the Cervix in PNG. Port Moresby: Government of Papua New Guinea; 2009.
7. Toliman P, Badman SG, Gabuzzi J, et al. Field evaluation of xpert HPV point-of-care test for detection of human papillomavirus infection by use of self-collected vaginal and clinician-collected cervical specimens. *J Clin Microbiol* 2016;54:1734-7.
8. Introducing and scaling up testing for human papillomavirus as part of a comprehensive programme for prevention and control of cervical cancer: a Step-by-step guide. Geneva: World Health Organization, 2020. Licence: CC BY-NC-SA 3.0 IGO
9. World Health Organization. (2019) WHO guidelines for the use of thermal ablation for cervical pre-cancer lesions. <https://apps.who.int/iris/handle/10665/329299>. Licence: CC BY-NC-SA 3.0 IGO

Elimination Implementation Checklist*

Strategic actions to achieve elimination targets

Completed

1	Develop a comprehensive costed National Cervical Cancer Elimination Strategy and seek endorsement among government, country leaders, policymakers, and communities	<input type="checkbox"/>
2	Communication and public outreach	
2.1	Develop a comprehensive communications strategy to accompany each component of the elimination strategy engaging health workers, community leaders, parents, teachers, and young people to maintain confidence in the programme and address stigma and misconceptions.	<input type="checkbox"/>
3	National HPV vaccination programme	<input type="checkbox"/>
3.1	Introduce the HPV vaccine for girls aged 9-14 years into national immunization programme.	<input type="checkbox"/>
3.2	Secure sufficient and affordable HPV vaccine and ensure an adequate cold-chain system for vaccine storage and delivery is in place.	<input type="checkbox"/>
3.3	Achieve and maintain high coverage of HPV vaccination by identifying appropriate multi-sectoral vaccination delivery platforms.	<input type="checkbox"/>
3.4	Establish or improve monitoring systems or vaccination registers to enable measurement of coverage and vaccine schedule adherence.	<input type="checkbox"/>
4	National cervical cancer screening and precancerous treatment programme	
4.1	Develop a national cervical cancer screening programme with clinical protocols for primary HPV testing and precancer treatment, involving relevant stakeholders when appropriate.	<input type="checkbox"/>
4.2	Integrate screening and precancer treatment into existing primary care and Universal Health Care (UHC) packages, including sexual and reproductive health services, HIV clinics, antenatal care.	<input type="checkbox"/>
4.3	Establish continuing professional development in-service programmes to build capacity of providers in cervical cancer screening and precancer treatment.	<input type="checkbox"/>
4.4	Understand social, financial, cultural, societal, and structural barriers to accessing services and create an enabling environment for cervical cancer screening and precancer treatment.	<input type="checkbox"/>
4.5	Strengthen laboratory capacity and quality assurance (QA) programmes and develop data systems that link laboratory information, screening registry data and other data systems (such as medical records and cancer registries).	<input type="checkbox"/>
5	Invasive cancer treatment and palliative care	
5.1	Develop and implement cervical cancer management guidelines and clinical protocols.	<input type="checkbox"/>
5.2	Establish effective referral pathways for women at all stages of care.	<input type="checkbox"/>
5.3	Strengthen pathology services, particularly at regional pathology centres and, if appropriate, make use of telepathology platforms to improve the capacity to interpret samples.	<input type="checkbox"/>
5.4	Expand surgical capacity through training programmes, and expand access to radiotherapy and chemotherapy services and strengthen oncology services.	<input type="checkbox"/>
5.5	Strengthen and integrate palliative care services by developing treatment plans that incorporate not only end-of-life care and pain relief, but also psychological and family support.	<input type="checkbox"/>
5.6	Optimize health workforce competencies throughout the continuum of care by establishing a long-term continuous training and education strategy for a multidisciplinary workforce.	<input type="checkbox"/>
5.7	Reduce cancer stigma by providing comprehensive support to enhance quality of life and address mental and sexual and reproductive health challenges faced by cancer survivors.	<input type="checkbox"/>
6	Monitoring and evaluation	
6.1	Strengthen governance and accountability of cervical cancer related programmes (HPV vaccination, cervical cancer screening, cancer treatment) and conduct regular reviews to ensure that national strategies, plans, and resource allocations reflect actual country needs.	<input type="checkbox"/>
6.2	Set country-specific targets, milestones, and indicators for monitoring and evaluating implementation of the National Cervical Cancer Elimination Strategy.	<input type="checkbox"/>
6.3	Improve current population-based cancer registries and develop new population-based cancer registries as needed to track the progress of the elimination targets.	<input type="checkbox"/>
6.4	Track patients throughout the continuum of services (screening, diagnosis and treatment).	<input type="checkbox"/>

*This checklist was developed based on the WHO strategy for cervical cancer elimination². For more guidance on the implementation of cervical cancer screening and treatment, see the WHO guide to strengthening cervical cancer prevention⁸ and the WHO guidelines for precancer treatment.⁹